

05/13/99
jc662 U.S. PTO

A/RE

PATENT/Docket No.: 4830.P RE

CERTIFICATE OF MAILING (37 CFR 1.10)

"Express Mail" No.: EM177638503 Date of Deposit: May 13, 1999

I hereby certify that this transmittal together with the patent application referred to below is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, DC 20231.

Julie K Lyons
Signature

Julie K. Lyons, Legal Assistant Name of Person Mailing Paper

jc542 U.S. PTO
05/13/99

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Commissioner of Patents and Trademarks
Washington, DC 20231

Sir:

Attached hereto for filing is an application for reissue of a U.S. Patent (hereafter the "original patent"), identified as follows:

Patent No. : 5,652,245
Issued Date : July 29, 1997
Inventor(s) : Arthur G. Romero
Title : Heterocyclic Amines Having Central Nervous System Activity

☒ This reissue application is being mailed by Express Mail under 37 CFR 1.10 and the required certificate appears above.

This reissue application includes the following parts as filed:

17 page(s) of specification
2 page(s) of claims
1 page(s) of abstract
0 sheets of drawing

ORDER FOR TITLE REPORT. Please prepare a certified Title Report (Abstract of Title) to the original patent. Title to the original patent is believed to be in the name of PHARMACIA & UPJOHN COMPANY, a Delaware corporation with a place of business in Kalamazoo, Michigan, by virtue of an assignment of the entire right and interest therein recorded at Reel 9719, Frame 0754.

The fee for preparing the Abstract of Title as required by 37 CFR 1.19(b)(2), may be charged to Deposit Account No. 21-0718. See the fee calculations below.

CONSENT OF ASSIGNEE. PHARMACIA & UPJOHN COMPANY, assignee of the original patent by virtue of the above-referenced assignment, hereby assents to the above-identified reissue application.

OFFER TO SURRENDER THE ORIGINAL LETTERS PATENT. PHARMACIA & UPJOHN COMPANY, through the undersigned, hereby offers to surrender the original patent upon grant of this application for reissue thereof in accordance with 37 CFR 1.178.

REISSUE DECLARATIONS. Attached hereto are:

1. ☒ a reissue declaration (37 CFR 1.63) and power of attorney by the assignee of the entire interest in the patent, inasmuch as the reissue application does not seek to enlarge the scope of the claims, or
☐ a reissue declaration by the inventor(s) (37 CFR 1.63) and power of attorney, inasmuch as the application may or may not seek to enlarge the scope of the claims, and
2. ☐ a statement under oath/declaration providing the particulars set forth in 37 CFR 1.175(a)(1)-(7) made by or on behalf of the applicant and, if applicable,
☐ corroborating statement(s) (37 CFR 1.175(b)).

FILING FEES. The filing fees for this reissue application are calculated as shown below:

	No. Filed	No. Extra	Rate	Fee
TOTAL CLAIMS FEE	12 - 20 =	0	x \$22	\$ 0.00
INDEPENDENT CLAIMS FEE	5 - 3 =	2	x \$80	\$ 160.00
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM			\$260	\$ 0.00
BASIC FEE				\$ 770.00
ABSTRACT OF TITLE REPORT			\$25	\$ 25.00
TOTAL FILING FEE				\$ 955.00

SPECIFIC DEPOSIT ACCOUNT AUTHORIZATION. Please charge my Deposit Account No. 21-0718 in the amount of the total filing fee above. Triplicate copies of this sheet are enclosed.

GENERAL DEPOSIT ACCOUNT AUTHORIZATION. The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or during the pendency of this reissue application or credit any overpayment to Deposit Account No. 21-0718.


- (1) Any additional filing fees or fees for the presentation of additional claims required under 37 CFR 1.16.
- (2) Any patent application processing fees under 37 CFR 1.17.

No authorization is given to charge the Issue Fee (37 CFR 1.18).

DELEGATION OF AUTHORITY. The authority of the undersigned to execute this document is derived from a delegation of authority from PHARMACIA & UPJOHN COMPANY, communicated to the Commissioner by letter dated January 27, 1999, a copy of which is attached to this paper.

Respectfully submitted,

Date: 11 May, 1999



Bruce Stein, Attorney
Registration No. 27,231
Telephone: (616) 833-1127

Mlg. Address: Intellectual Property Legal Services, Pharmacia & Upjohn Company, Kalamazoo, MI 49001

Attachments: ☒ Delegation of authority
 ☒ Application (including specification, claims, abstract and drawings, if any)
 ☒ Declaration, including Power of Attorney and Statement under 37 CFR 1.175(a), by ASSIGNEE
 ☐ Corroborating statement(s) under 37 CFR 1.175(b)

Enclosure: Return postcard

CERTIFICATE UNDER 37 C.F.R. 3.73(b)

Applicant: Arthur G. Romero

Application No.: 08/592,328

Filed: 1-23-96

For: Heterocyclic Amines Having Central Nervous System Activity

Pharmacia & Upjohn Company, a

(Name of Assignee)

Corporation

(Type of Assignee, e.g. corporation, partnership, university, government agency, etc.)

certifies that it is the assignee of the entire right, title and interest in the patent application identified above by virtue of either:

A. ☒ An assignment from the inventor(s) of the patent application identified above. The assignment was recorded in the Patent and Trademark Office at Reel 9719, Frame 0754, or for which a copy thereof is attached.

OR

B. ☐ A chain of title from the inventor(s), of the patent application identified above, to the current assignee as shown below:

1. From: _____ To: _____
The document was recorded in the Patent and Trademark Office at Reel _____, Frame _____, or for which a copy thereof is attached.

2. From: _____ To: _____
The document was recorded in the Patent and Trademark Office at Reel _____, Frame _____, or for which a copy thereof is attached.

3. From: _____ To: _____
The document was recorded in the Patent and Trademark Office at Reel _____, Frame _____, or for which a copy thereof is attached.

☐ Additional documents in the chain of title are listed on a supplemental sheet.

☒ Copies of assignments or other documents in the chain of title are attached.

The undersigned has reviewed all the documents in the chain of title of the patent application identified above and, to the best of undersigned's knowledge and belief, title is in the assignee identified above.

The undersigned (whose title is supplied below) is empowered to act on behalf of the assignee.

I hereby declare that all statements made herein of my own knowledge are true, and that all statements made on information and belief are believed to be true; and further, that these statements are made with the knowledge that willful false statements, and the like so made, are punishable by fine or imprisonment, or both, under Section 1001, Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date : 12 May 1999

Name : Bruce Stein

Title : Patent Attorney

Signature: Bruce Stein

HETEROCYCLIC AMINES HAVING CENTRAL NERVOUS SYSTEM ACTIVITY

The present patent application is a national phase of International application No. PCT/US94/06648. International filing date 17 Jun. 1994, which was a continuation in part of U.S. patent application Ser. No. 08/097,608, filed 27 Jul. 1993, now abandoned.

BACKGROUND OF THE INVENTION

The present invention is directed toward tricyclic nitrogen containing compounds, heterocyclic amines, having anxiolytic and anti-depressant activity. These new compounds are suitable for treating central nervous system disorders including schizophrenia, Parkinson's disease, anxiety or as compounds for lowering blood pressure or treating migraines.

A series of dihydropbenzalenes, tricyclic amine substituted compounds, and related compounds having central nervous system activity were described in PCT Int. Pub. No. WO87/04153 and in PCT Int. Pub. No. WO88/04292. A major difference between those compounds and the present invention is that the subject compounds have at least one nitrogen atom in the tricyclic ring structure which is shared by two of the ring structures. Generally, the subject compounds have exhibited anxiolytic activity and better oral bioavailability.

INFORMATION DISCLOSURE STATEMENT

PCT Int. Pub. No. WO87/04153 and PCT Int. Pub. No. WO88/04292 each describe tricyclic structures having central nervous system activity.


U.S. Pat. No. 4,110,339 discloses 4-(di-n-propyl)amino-1,3,4,5-tetrahydrobenz(cd)indole compounds useful as prolactin inhibitors and in the treatment of Parkinsonism. European Patent Application 153,083 and German Patent 3,346,573 disclose methoxy substituted 4-(di-n-propyl)amino-1,3,4,5-tetrahydrobenz(cd)indole compounds. These publications disclose nitrogen containing tricyclic ring structures but the nitrogen is not shared by any of the rings.

Evans, D. D., Peters, D. J., *J. Chem. Soc., Perkin Trans. 1*, pp 285-88 (1974) discloses nitrogen containing tricyclic ring structures where the nitrogen is shared by two ring structures but additionally includes other substituents not common to the subject compounds.

PCT Int. Pub. No. WO 90/15058 discloses compounds with the characteristic tricyclic nitrogen containing structure of the subject invention with the exception of the ring nitrogen "X" substituents.

SUMMARY OF THE INVENTION

In one aspect, the present invention is directed toward tricyclic nitrogen containing compounds of Formula I:



and pharmaceutically acceptable salts thereof. R₁ and R₂ are independently hydrogen, C₁₋₆ alkyl or R₁ and R₂ are joined to form pyrrolidine, piperidine, morpholine or imidazole. X

is OCH_3 , SO_2R_3 , SO_2CF_3 or CN where R_3 is C_{1-6} alkyl or an Aryl; and Y is hydrogen, Cl , Br , F , CN , CONR_1R_2 , CF_3 , OCH_3 , $\text{SO}_2\text{NR}_1\text{R}_2$.

These new compounds have been found to exhibit anxiolytic activity in isolation induced aggression and plasma corticosterone models. They are also suitable for treating various central nervous system disorders effected by 5-HT_{1A} receptors such as, schizophrenia, Parkinson's disease, anxiety, depression, or as compounds for lowering blood pressure and treating migraine headaches in patients in need of such treatments.

In yet another aspect, the present invention is a method for treating central nervous system (CNS) disorders influenced by 5-HT_{1A} receptors such as anxiety, depression, hypertension and associated high blood pressure, Parkinson's disease and schizophrenia in animal or human hosts by administering a pharmaceutically effective amount of a compound of Formula I including pharmaceutically acceptable salts. Other uses for these compounds include panic attacks, eating disorders, obsessive-compulsive disturbances seen in dementia disorders. In addition, central 5-HT receptor activation are believed to be involved in mediating sexual behavior and therefore these compounds would be useful to stimulate sexual activity and to alleviate impotence.

DETAILED DESCRIPTION OF THE INVENTION

The present invention describes compounds of Formula I, above, having central nervous system activity. The compounds are characterized by a tricyclic ring structure having a shared nitrogen atom between two rings, an amine substituent (NR_1R_2) and a substituted ring nitrogen (X) as structurally depicted by Formula I. The systematic names for the ring systems in these compounds may be found by consulting the Ring Systems Handbook, 1988 edition, published by Chemical Abstracts Service. These names are derived by combining the names of benzene or a monocyclic heterocycle with the name of a bicyclic heterocycle to which it is fused. The atoms and bonds common to the fused rings are then specified to distinguish it from isomeric systems with similar names.

The particular compounds have been found to be active in various central nervous system screens such as hypothermia and hypoxic stress tests and have been found to be dopamine and serotonin such as, 5HT_{1A} receptor binding assay antagonists.

The following definitions are applied to the structural formula represented by Formula I, above.

" $\text{C}_1\text{-C}_6$ alkyl" means methyl, ethyl, propyl, butyl, pentyl and hexyl and isomeric forms thereof.

"Aryl" means aromatic ring structures containing five to ten carbon atoms which can be optionally substituted with halogen atoms, C_{1-6} alkyl (which can be optionally substituted with halogen and hydroxyl groups) and hydroxyl groups such as phenyl, α -naphthyl, β -naphthyl, *m*-methylphenyl, *p*-trifluoromethylphenyl and the like. Aryl also includes the various heteroaryl groups which contain the heteroatoms nitrogen, sulfur or oxygen to form pyridine, thiophene, furan, pyrimidine, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 3-pyrazinyl, 2-quinolyl, 3-quinolyl, 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 2-quinazolinyl, 4-quinazolinyl, 2-quinoxaliny, 1-phthalaziny, 2-imidazolyl, 4-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl,

4-thiazolyl, 5-thiazolyl, 2-indolyl, 3-indolyl, 3-indazolyl, 2-benzoxazolyl, 2-benzothiazolyl, 2-benzimidazolyl, 2-benzofuranyl, 3-benzofuranyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,4-triazol-3-yl, 1,2,4-triazol-5-yl, 1,2,3,4-tetrazol-5-yl, 5-oxazolyl, 1-pyrrolyl, 1-pyrazolyl, 1,2,3-triazol-1-yl, 1,2,4-triazol-1-yl, tetrazolyl, 1-indolyl, 1-indazolyl, 2-isoindolyl, 1-purinyl, 3-isothiazolyl, 4-isothiazolyl and 5-isothiazolyl.

"Pharmaceutically acceptable salts" are hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate, acetate, propionate, lactate, maleate, malate, succinate, tartrate, cyclohexanesulfamates, methanesulfonates, ethanesulfonates, benzenesulfonates, toluenesulfonates and other pharmaceutically acceptable counter ions for amines. Additionally, the compounds of this invention may be administered in a suitable hydrated form.

The compounds of the invention include both racemic and optically pure products which can be separated by conventional methods into the R- and S-isomers. Resolution can be accomplished using resolving agents such as optically active dibenzoyltartaric acid, camphorsulfonic acid, bis-o-toluoityltaric acid, tartaric acid, and diacetyl tartaric acid.

A second procedure useful in resolving primary and secondary amine compounds of Formula I involves their conversion to diastereomeric amides using an optically active acid. The diastereomeric amides are separated and the amide bond is cleaved to afford the optically pure Formula I compounds. This procedure is illustrated in PCT International Publication No. WO 90/15058 (Examples 49 and 50) for the preparation of the optical isomers using t-butoxycarbonyl-L-phenylalanine as the resolving agent. For the resolution, the racemic compound is coupled to t-butoxycarbonyl-L-phenylalanine and the diastereomeric amide products are separated by chromatography into the (+) and (-) forms. The (-) isomer is reacted with trifluoroacetic acid to give (-) N-(5,6-dihydro-2-oxo-4H-imidazo(4,5,1-ij)quinolin-5-yl)-L-phenylalanineamide. Edman degradation of this compound, by reaction with phenyl isothiocyanate followed by trifluoroacetic acid, removes the phenylalanine residue and affords the (-) form of the compound. Further reaction of this product with propionaldehyde and sodium cyanoborohydride gives the (-) form of the active isomer.

General procedures for preparing compounds of Formula I are shown in Schemes 1 and 2, below, and as cross-referenced and described in the Examples. Methods for preparing various intermediates for the subject compounds are described in PCT International Publication No. WO 90/15058 herein incorporated by reference.

The dosage regimen for treating patients with the compounds of this invention is selected in accordance with a variety of factors including the type, age, weight, sex, and medical condition of the patient, the severity of the psychosis, the route of administration and the particular compound employed. An ordinarily skilled physician or psychiatrist will readily determine and prescribe the effective amount of compound to prevent or arrest the progress of the condition. In so proceeding, the physician or psychiatrist could employ relatively low dosages at first, subsequently increasing the dose until a maximum response is obtained.

Initial dosages of the compounds of the invention are ordinarily in the area of at least 10 mg up to about 1200 mg

per day orally, which may be given in a single dose or in multiple doses. When other forms of administration are employed equivalent doses are administered. When dosages beyond 600 mg are employed, care should be taken with each subsequent dose to monitor possible toxic effects.

The compounds of this invention are administered in oral unit dosage forms such as tablets, capsules, pills, powders, or granules. They may also be introduced parenterally, (e.g., subcutaneously, intravenously, or intramuscularly), using forms known to the pharmaceutical art. They also may be administered rectally or vaginally in such forms as suppositories or bougies. In general, the preferred route of administration is oral.

The compounds of this invention can also be administered as pharmaceutically or therapeutically acceptable salt such as hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate, acetate, propionate, lactate, maleate, malate, succinate, tartrate, cyclohexanesulfamates, methanesulfonates, ethanesulfonates, benzenesulfonates, toluenesulfonates and the like. Additionally, the compounds of this invention may be administered in a suitable hydrated form.

Compounds of the subject invention were evaluated in an isolation-induced aggression assay to measure their ability to control or arrest aggressive behavior as measured against a saline "Control". Male CF-1 mice (Charles River Labs) were housed singly in wire cage drawers for several weeks of isolation and aggression training. After this, the isolated mouse (weighing 30–50 g) initiated an "attack" whenever an intruder mouse was placed into his cage. Intruders were male CF-1 Charles River mice, about the same size as isolated mice and housed in groups of 12 per cage. For drug testing, the isolated mice were dosed orally (p.o.) with drug or 0.25% methylcellulose vehicle. Thirty minutes later, an intruder mouse was introduced and the latency to attack was recorded. Treatment groups (n=6), including a positive control, were tested by an observer blinded to the treatments, and there was an additional non-blinded vehicle group. The results were as follows:

Compound	Seconds
Control	66
Example 1 ¹	600
Control	152
Example 2 ²	565
Control	66
Example 2 ³	600

¹Formula I where R₁ is hydrogen, R₂ is methyl, Y is hydrogen and X is —OCH₃ (R-enantiomer);

²Formula I where R₁ is methyl, R₂ is methyl, Y is hydrogen and X is —OCH₃ (racemate);

³Formula I where R₁ is methyl, R₂ is methyl, Y is hydrogen and X is —OCH₃ (R-enantiomer).

Compounds of Example 4 (Formula I where R₁ and R₂ are propyl, Y is hydrogen and X is —OCH₃) were evaluated in an receptor binding assay (5HT Ligand) for calculation of IC₅₀ values and K_i (nM) values. Receptor binding assays are well known tests for evaluating compounds for activity as described in "Cloning and Pharmacological Characterization of a Novel Human 5-hydroxy-1-tryptamine_{1D} Receptor Sub-Type", Mol. Pharm., 42 439–44 (1992), herein incorporated by reference. The results were as follows:

Compounds Ex. 4	Receptor	Ligand	Ki (nM)
Racemate	5-HT1DA-Clone	5HT	15.30
"	5-HT1DB-Clone	5HT	7.17
R-Enantiomer	5-HT1DA-Clone	5HT	38.00
"	5-HT1DB-Clone	5HT	58.00
"	5-HT1A-Clone	DPAT	97
S-Enantiomer	5-HT1A-Clone	DPAT	904

The above data shows that the racemate compound is selective for the 5HT_{1DB} receptor and that the activity resides in the R-enantiomer compound.

EXAMPLE 1

(R)-5-Methylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (8a)

Preparation of Methyl (R)-N-(1,2,3,4-tetrahydro-1-methoxy-2-oxo-3-quinolinyl)carbamate (2a):

A solution of methyl chloroformate (42.8 g, 0.453 mol) in chloroform (50 ml) was added to a mixture of (R)-1,2,3,4-tetrahydro-1-methoxy-2-oxo-3-quinolinamine 1 (M. Kawase, T. Kitamura, Y. Kikugawa, J. Org. Chem., 54, 1989 3394-3403) (0.15 mol) and sodium bicarbonate (50.4 g, 0.600 mol) in chloroform (250 ml) and water (100 ml) over a period of 10 minutes at 0° C. The mixture was stirred overnight at room temperature, diluted with pentane, and the layers were separated. The aqueous layer was extracted with diethylether, and the combined organic extracts were washed with brine and dried (MgSO₄). The solvent was removed under vacuum to leave an amber oil. Purification by flash chromatography (230-400 mesh silica gel, 35-40% ethyl acetate in hexane) gave the title compound as a light yellow oil (37.0 g, 99%).

NMR (CDCl₃, TMS) δ 2.865 (t, t_J =14.7 Hz, 1H), 3.437 (dd, J =6.0 & 15.3 Hz, 1H), 3.723 (s, 3H, NOCH₃), 3.932 (s, 3H, COOCH₃), 4.426 (dt, J =5.8 & 14.2 Hz, 1H, N—CH), 5.856 (br. s, 1H, NH), 7.098 (t of d, J =1.2 & 7.4 Hz, 1H), 7.228 (m, 2H), 7.331 (t, t_J =7.7 Hz, 1H).

$[\alpha]_D^{25}$ =-47° (25° C., CHCl₃, c =0.9977).

Preparation of (R)-3-Methylamino-1,2,3,4-tetrahydroquinoline (3a):

A solution of (R)-N-(1,2,3,4-tetrahydro-1-methoxy-2-oxo-3-quinolinyl)carbamate (2a, 36.25 g, 0.145 mol) in tetrahydrofuran (400 ml) was cooled in ice, and borane dimethylsulfide complex (10M, 73 ml, 0.73 mol) was added. The ice bath was removed, and mixture slowly came to a vigorous reflux which required cooling with an ice bath. When the reaction subsided, it was heated at reflux on the steam bath overnight. The mixture was cooled in ice, and water (80 ml) and then 10% aqueous hydrochloric acid (70 ml) were added dropwise. The mixture (acidic by pH paper) was stirred at reflux on the steam bath for 1.25 hr. The mixture was cooled in ice and basified slowly with solid sodium hydroxide. The mixture was diluted with water and pentane, and the layers were separated. The aqueous was extracted with diethylether, and the combined organic extracts were dried (MgSO₄). The solvent was removed under vacuum to leave an oil (23.96 g, 100%). A sample (1.3 g) was purified via gravity chromatography (70-230 mesh silica gel; 6% methanol, 0.6% ammonium hydroxide, chloroform) to give the title compound as a light yellow oil (0.84 g).

NMR (CDCl₃, TMS) δ 1.529 (br. s, 1H, methylamine NH), 2.503 (s, 3H, N—CH₃), 2.62-2.71 (m, 1H), 2.93-3.14 (m, 3H), 3.394 (d of t, J =2.2 & 10.9 Hz, 1H), 3.814 (br.

s.1H,NH), 6.483 (d, $J=7.7$ Hz, 1H), 6.624 (d of t, $J=1.0$ & 7.4 Hz, 1H), 6.95–6.98 (m, 2H).

The compound (0.83 g, 5.12 mmol) was dissolved in methanol and combined with a solution of trimethylsilyl chloride (0.54 g, 5.0 mmol) in methanol, and the solvent was removed under vacuum. The residue was crystallized from methanol/diethylether to give the hydrochloride salt of the title compound (3a) as light yellow-green crystals (0.56 g, m.p. 193°–195° C.).

10 $[\alpha]_D^{25} = +26^\circ$ (25° C., CH₃OH, $c=1.0232$).

Preparation of (R)-N-Methyl-N-[3-((1-trifluoromethylacetyl)-1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide (4a)

A solution of (R)-3-Methylamino-1,2,3,4-tetrahydroquinoline (3a, 21.32 g, 0.131 mol) in tetrahydrofuran (200 ml) was cooled in ice, and trifluoroacetic anhydride (69.2 g, 0.329 mol) was added dropwise. The mixture was stirred at 0° C. for 1 hr and at room temperature for 2 hr. The mixture was again cooled in ice and water (150 ml) was added. Solid sodium bicarbonate was added portionwise until the evolution of gas ceased. The mixture was extracted three times with diethylether, and the combined organic extracts were washed with saturated sodium bicarbonate and brine. The solution was dried (MgSO₄), and the solvent was removed under vacuum to leave a yellow oil (42.2 g). Purification by flash chromatography (230–400 mesh silica gel; 15% ethyl acetate in hexane) gave the title compound as a yellow oil (31.12 g, 67% yield).

$[\alpha]_D^{25} = +59^\circ$ (25° C., CHCl₃, $c=0.5495$).

10 MR (CDCl₃, TMS) δ 2.75–3.40 (m, 5H), 3.65–3.95 (m, 1H), 3.95–4.30 (m, 1H), 4.572 (quintet, $J=7.6$ Hz, 0.3H), 4.921 (m, 0.7H), 7.15–7.45 (m, 3.7H), 7.697 (m, 0.3H).

Preparation of (R)-3-Methylamino-1-trifluoromethylacetyl-1,2,3,4-tetrahydroquinoline (5a):

5 A solution of (R)-N-Methyl-N-[3-((1-trifluoromethylacetyl)-1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide (4a, 29.78 g, 84.1 mmol) in tetrahydrofuran (140 ml) was cooled in ice and a solution of potassium hydroxide (45.6% solution, 7.5 ml, 89.3 mmol) in water (40 ml) was added. The mixture was stirred at 0° C. for 30 minutes and at room temperature for 1 hour. The mixture was extracted twice with diethylether. The combined organic extracts were washed with brine and dried (MgSO₄). The solvent was removed under vacuum to leave a yellow oil (21.7 g, 100%). A sample (1.05 g) was purified via flash chromatography (230–400 mesh silica gel, 15% ethyl acetate in hexane) gave a yellow oil (0.90 g). Crystallization from ethyl acetate/hexane gave the title compound as a colorless solid (0.45 g, m.p. 68°–69° C.).

15 NMR (CDCl₃, TMS) δ 2.871–3.182 (m, 5H, N—CH₃ & Ar—CH₂), 3.33–3.449 (m, 2H, N—CH₂), 3.890 (br. s, 1H, NH), 4.35–4.48 (m, 0.4H, N—CH), 4.800–4.888 (m, 0.6H, N—CH), 6.553 (t, $J=7.6$ Hz, 1H), 6.660–6.735 (m, 1H), 6.996–7.068 (m, 2H).

$[\alpha]_D^{25} = -29^\circ$ (25° C., CHCl₃, $c=0.9917$).

Preparation of (R)-N-Methyl-N-[3-(1-(N-methoxyaminocarbonyl)-1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide (6a):

A solution of phosgene (1.93M in toluene, 15.6 ml, 30.1 mmol) in dry tetrahydrofuran (75 ml) was cooled in ice and a solution of (R)-3-Methylamino-1-trifluoromethylacetyl-1,2,3,4-tetrahydroquinoline (5a, 7.75 g, 30.0 mmol) and triethylamine (4.20 ml, 30.1 mmol) in dry tetrahydrofuran (75 ml) was added dropwise. The mixture was stirred in an ice bath for 50 minutes, and methoxylamine hydrochloride (5.01 g, 60.0 mmol) and triethylamine (8.40 ml, 60 mmol) were added. The mixture was stirred at room temperature for

24 hours, diluted with diethylether, and washed with water, 10% hydrochloric acid, saturated sodium bicarbonate solution, and brine. The solution was dried (MgSO_4), and the solvent was removed under vacuum to leave a oil (8.85 g). Purification by flash chromatography (230–400 mesh silica gel, 50% ethyl acetate in hexane) gave the title compound as a colorless oil (7.42 g, 75% yield).

NMR (CDCl_3 , TMS) δ 2.817–3.097 (m, 5H, N—CH₃ & Ar—CH₂), 3.66–4.035 (m, 5H, N—CH₂ & O—CH₃), 4.38–4.863 (m, 1H, N—CH), 7.148–7.366 (m, 4H), 7.813 & 7.829 (two s, 1H, NH).

$[\alpha]_D^{25} +5^\circ$ (25° C., CHCl_3 , $c=1.137$).

Preparation of (R)-5-N-(N-Methyltrifluoromethylacetamido)-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (7a):

A solution of (R)-N-Methyl-N-[3-(1-(N-methoxyaminocarbonyl)-1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide 6a, 6.61 g, 20.0 mmol) in chloroform (100 ml) was degassed with argon and bis (trifluoroacetoxy)iodobenzene (10.32 g, 24 mmol) was added portionwise. The solution was stirred at reflux for 7 minutes. The reaction mixture was cooled, diluted with pentane, and washed twice with saturated sodium bicarbonate solution and once with brine. The solution was dried (MgSO_4), and the solvent was removed under vacuum to leave a red-brown oil (11.5 g). Purification by flash chromatography (230–400 mesh silica gel; 35–40% ethyl acetate in hexane) gave an amber solid (4.74 g, 72% yield). Crystallization from ethyl acetate/hexane gave the title compound as off-white crystals (4.55 g, m.p. 148.5°–150.5° C.).

NMR (CDCl_3 , TMS) δ 2.995–3.299 (m, 5H, N—CH₃ & Ar—CH₂), 3.724–3.839 (m, 1H), 4.089 & 4.099 (two s, 3H, O—CH₃), 5.059–4.227 (m, 1H), 4.520 (heptet, $j=5.4$ Hz, 0.3H), 4.931 (heptet, $j=5.1$ Hz, 0.7H), 6.900–7.121 (m, 3H).

$[\alpha]_D^{25} +38^\circ$ (25° C., CHCl_3 , $c=1.0215$).

Preparation of (R)-5-Methylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (8a):

Potassium carbonate (2.05 g, 14.8 mmol) was added to a solution of (R)-5-N-(N-methyltrifluoromethylacetamido)-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (7a, 3.75 g, 11.4 mmol) in methanol (75 ml). The mixture was stirred at reflux for 7 hours. The solvent was removed under vacuum, and the residue was partitioned between diethylether and water (75 ml). The aqueous solution was saturated with brine and extracted twice more with diethylether. The combined organic extracts were dried (MgSO_4), and the solvent was removed under vacuum to leave an oil. After standing for three days, the aqueous developed a precipitate which was extracted twice with 75% tetrahydrofuran in diethylether. The combined organic extracts were dried (MgSO_4), and the solvent was removed under vacuum to leave a brown semisolid. The combined samples were purified via flash chromatography (230–400 mesh silica gel, 70–80% tetrahydrofuran in ethyl acetate to give the title compound as an amber oil (2.7 g, 100%).

NMR (CDCl_3 , TMS) δ 2.545 (s, 3H, N—CH₃), 2.781 (dd, $j=7.4$ & 16.1 Hz, 1H), 3.071 (dd, $j=4.0$ & 16.0 Hz, 1H), 3.233 (heptet, $j=3.8$ Hz, 1H), 3.645 (dd, $j=7.0$ & 12.4 Hz, 1H), 4.045 (m, 1H), 4.088 (s, 3H, NOCH₃), 6.893 (d, $j=7.4$ Hz, 1H), 6.964 (d, $j=7.1$ Hz, 1H), 7.039 (t, $j=7.7$ Hz, 1H).

A sample (0.70 g) was dissolved in tetrahydrofuran, diluted with diethyl ether, and excess ethereal hydrochloric acid was added. The precipitate was filtered, washed with diethylether, and crystallized from methanol/diethylether to give the hydrochloride salt of the title compound (8a) as tan crystals (0.68 g, m.p. 196°–197° C.).

$[\alpha]_D^{25} -26^\circ$ (25° C., CH_3OH , $c=1.0139$).

EXAMPLE 2

(R)-5-Dimethylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (9a)

Following the procedure of Example 1, a solution of (R)-5-Methylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,

5.1-ij]quinolin-2(1H)-one (8a, 1.88 g, 8.06 mmol) in methanol (40 ml) was stirred at room temperature, and 37% aqueous formaldehyde (2.40 ml, 32 mmol) and acetic acid (0.98 g, 16.2 mmol) were added. The mixture was cooled in ice, and sodium cyanoborohydride (1.15 g, 17.0 mmol) was added portionwise over a 5 minute period. The mixture was stirred at 0° C. for 5 minutes and at room temperature for 6 hours. The solvent was removed under vacuum, and the residue was diluted with 10% sodium carbonate solution and extracted twice with diethylether. The combined organic extracts were washed with brine and dried (MgSO₄). The solvent was removed under vacuum to leave an amber oil (2.06 g). Purification by flash chromatography (230–400 mesh silica gel, 65% tetrahydrofuran in ethyl acetate) gave the rifle compound as an amber oil (1.77 g, 89% yield).

NMR (CDCl₃, TMS) δ 2.436 (s, 6H, N—CH₃), 2.853–3.068 (m, 3H), 3.525–3.596 (m, 1H), 4.081 (s, 3H, O—CH₃), 4.188–4.243 (m, 1H), 6.890 (d, J=7.4 Hz, 1H), 6.954 (d, J=7.1 Hz, 1H), 7.033 (t, J=7.6 Hz, 1H).

The compound was dissolved in diethylether and excess ethereal hydrochloric acid was added. The precipitate was filtered, washed with ether and crystallized from methanol/diethylether to give the hydrochloride salt of the title compound (9a) as yellow crystals (1.76 g, 195° C.).

[α]_D²⁰ = –19° (25° C., CH₃OH, c=1.0234).

EXAMPLE 3

(R)-5-Propylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (8b)

(R)-N-(1,2,3,4-Tetrahydro-1-methoxy-2-oxo-3-quinolinyl)propanamide (2b):

The amine (1, 33.51 g, 0.135 mol) was dissolved in propionic anhydride (78.2 g, 0.60 mol) while controlling the temperature with an ice bath. The mixture was stirred at room temperature for 5 hours, and water (100 ml) was added, saturated NaHCO₃ (200 ml) was slowly added, and then 10% Na₂CO₃ was slowly added to neutralize the mixture to pH 7.5. The mixture was extracted with diethylether, saturated with sodium chloride, and extracted again with diethylether and a mixture of diethylether and chloroform. The combined extracts were washed with 5% Na₂CO₃ and brine. The solution was dried (MgSO₄), and the solvent was removed under vacuum to leave an amber oil. Purification by flash chromatography (230–400 mesh silica gel, 60% ethyl acetate in hexane) gave an oil. A sample was crystallized from ethyl acetate/hexane to give the title compound (2b) as colorless crystals (m.p. 98°–99° C.).

[α]_D²⁰ = –100° (25° C., CHCl₃, c=0.9698).

NMR (CDCl₃, TMS) δ 1.206 (t, J=7.6 Hz, 3H, CO—CH₃), 2.343 (q, J=7.6 Hz, 2H, COCH₂), 2.768 (t, J=14.7 Hz, 1H), 3.527 (dd, J=6.1 & 15.1 Hz, 1H), 3.934 (s, 3H, O—CH₃), 4.614 (d of t, J=5.6 & 14.2 Hz, 1H, N—CH), 6.633 (br, d, J=4.5 Hz, 1H, NH), 7.099 (d of t, J=1.2 & 7.4 Hz, 1H), 7.230 (d, J=7.9 Hz, 2H), 7.327 (t, J=7.7 Hz, 1H).

Preparation of (R)-1,2,3,4-Tetrahydro-N-propyl-3-quinolinamine monohydrochloride (3b):

Borane methyl sulfide (10M, 42.5 ml, 0.425 mol) was added to a solution of (R)-N-(1,2,3,4-Tetrahydro-1-methoxy-2-oxo-3-quinolinyl)propanamide (2b, 21.29 g, 85.7 mmol) in tetrahydrofuran (250 ml) with stirring at room temperature. The reaction was exothermic, and an ice bath was used to control the reaction. The mixture was stirred at 0° C. for 30 minutes, at room temperature overnight, and then at reflux for 4 hours. The mixture was cooled in ice, and water (50 ml) was carefully added dropwise. When the

addition was complete. 10% HCl was added dropwise until the mixture was acidic by pH paper. The mixture was stirred at room temperature for 1 hour and at reflux for 1 hour. The mixture was washed with pentane and filtered. The filtrate was basified with 15% sodium hydroxide and extracted with diethylether. The aqueous layer was saturated with sodium chloride and extracted twice with diethylether. The combined ether extracts were dried (MgSO_4), and the solvent was removed under vacuum to leave an oil (14.8 g, 91%). A sample (1.0 g) was purified via gravity chromatography (70–230 mesh silica gel, 90 g; 3% methanol, 0.3% NH_4OH , CHCl_3) to give the title compound as a colorless oil.

NMR (CDCl_3 , TMS) δ 0.924 (t, $J=7.4$ Hz, 3H, CH_3), 1.518 (sextet, $J=7.4$ Hz, 3H), 2.641–2.711 (m, 3H), 2.965 (dd, $J=4.0$ & 14.7 Hz, 1H), 3.048–3.131 (m, 2H), 3.361–3.432 (m, 1H), 3.825 (br. s, 1H), 6.488 (d, $J=7.9$ Hz, 1H), 6.626 (d of t, $J=1.0$ & 7.4 Hz, 1H), 6.951–7.002 (m, 2H).

The compound (0.85 g, 4.47 mmol) was combined with a solution of trimethylsilyl chloride (0.57 ml, 4.49 mmol) in methanol. The solvent was removed under vacuum to leave a solid, which was crystallized twice from methanol/diethylether to give the hydrochloride salt of the title compound (3b) as off-white crystals (m.p. $185^\circ\text{--}192^\circ\text{C}$). $[\alpha]_D^{25} = +12^\circ$ (CH_3OH , 25°C , $c=0.8073$).

NMR (CD_3OD , TMS) δ 1.038 (t, $J=7.4$ Hz, 3H, CH_3), 1.705–1.833 (m, 2H, propyl N—C— CH_2), 2.986 (dd, $J=5.5$ & 17.2 Hz, 1H), 3.065–3.120 (m, 2H, N— CH_2), 3.265–3.342 (m, 1H), 3.436–3.572 (m, 2H), 3.734–3.800 (m, 1H), 6.731–6.805 (m, 2H), 7.037–7.084 (m, 2H).

Preparation of (R)-N-Propyl-N-[3-((1-trifluoromethylacetyl)-1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide (4b):

A solution of (R)-1,2,3,4-tetrahydro-N-propyl-3-quinolinamine (3b, 14.73 g, 77.4 mmol) in tetrahydrofuran (50 ml) was cooled in ice and trifluoroacetic anhydride (40.6 g, 0.193 mol) was added dropwise. When the addition was complete, the cold bath was removed, and the mixture was stirred overnight at room temperature. Water (75 ml) was added, and sodium bicarbonate was added portionwise until the mixture was basic to litmus. The mixture was extracted three times with diethylether, and the combined extracts were washed with saturated sodium bicarbonate solution and brine. The solution was dried (MgSO_4), and the solvent was removed under vacuum to leave an amber oil (28.4 g). A sample (0.87 g) was purified via flash chromatography (230–400 mesh silica gel, 5–6% ethyl acetate in hexane) to give a colorless oil which was crystallized from hexane to give the title compound (4b) as colorless crystals (0.61 g, m.p. $77^\circ\text{--}79^\circ\text{C}$).

NMR (CDCl_3 , TMS) δ 0.897–0.985 (m, 3H, propyl CH_3), 1.706 (sextet, $J=7.7$ Hz, 2H, propyl N—C— CH_2), 2.85–3.69 (m, 4H), 3.69–4.38 (m, 2.6H), 4.500 (quintet, 0.4H), 7.15–7.40 (m, 3.6H), 7.726 (br. s, 0.4H).

$[\alpha]_D^{25} = +8^\circ$ (25°C , CHCl_3 , $c=1.0067$).

Preparation of (R)-N-Propyl-N-[3-((1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide (5b):

A solution of N-Propyl-N-[3-((1-trifluoromethylacetyl)-1,2,3,4-tetrahydroquinolinyl)]trifluoromethylacetamide (4b, 27.64 g, 72.3 mmol) in tetrahydrofuran (200 ml) was cooled in ice, and a solution of potassium hydroxide (45.6% in water, 6.07 ml, 72.3 mmol) was added. The mixture was stirred at 0°C for 15 minutes and at room temperature for 30 minutes. The mixture was diluted with diethylether, and the layers were separated. The aqueous layer was extracted with diethylether, and the combined organic extracts were washed with brine and dried (MgSO_4). The solvent was removed under vacuum to leave an oil (21.4 g). Purification

by flash chromatography (230–400 mesh silica gel; 10% ethyl acetate in hexane) gave a yellow oil (16.68 g, 81% yield). A sample (0.75 g) was rechromatographed using the same conditions to give an oil which crystallized (0.66 g).
 5 Crystallization from hexane gave the title compound (5b) as colorless crystals (0.54 g; m.p. 68°–69° C.).

NMR (CDCl₃, TMS) δ: 0.883–0.946 (m, 3H, propyl CH₃), 1.58–1.77 (m, 2H, propyl N—C—CH₂), 2.83–2.95 (m, 1H), 3.104–3.483 (m, 4.5H), 3.708 (t, J=10.4 Hz, 0.5H), 3.893 (br.
 10 s, NH), 4.16–4.28 & 4.32–4.46 (m, 1H, N—CH), 6.546 (t, J=6.9 Hz, 1H), 6.647–6.725 (m, 1H), 6.965–7.063 (m, 2H).

[α]_D²⁵=62° (25° C., CH₃OH, c=0.9991).

Preparation of (R)-N-Propyl-N-[3-(1-(N-methoxyaminocarbonyl)-1,2,3,4-tetrahydroquinolinyl)]
 15 trifluoromethylacetamide (6b):

A solution of phosgene (1.93M in toluene, 10.4 ml, 20.0 mmol) in dry tetrahydrofuran (50 ml) was cooled in ice and a solution of N-Propyl-N-[3-(1,2,3,4-tetrahydroquinolinyl)] trifluoromethylacetamide (5b, 5.73 g, 20.0 mmol) and triethylamine (2.03 g, 20.1 mmol) in dry tetrahydrofuran (50 ml) was added over 5 minutes. The mixture was stirred in ice for 30 minutes, and methoxylamine hydrochloride (3.34 g, 40.0 mmol) and triethylamine (4.06 g, 40.2 mmol) were added. The mixture was stirred at room temperature for 24
 25 hours, diluted with diethylether and washed with water, twice with 10% hydrochloric acid solution, saturated sodium bicarbonate solution and brine. The solution was dried (MgSO₄), and the solvent was removed under vacuum to leave an oil (7.88 g). Purification by flash chromatography
 30 (230–400 mesh silica gel; 40% ethyl acetate in hexane) gave the title compound (6b) as an oil (6.25 g).

NMR (CDCl₃, TMS) δ: 0.899–0.969 (m, 3H, propyl CH₃), 1.62–1.83 (m, 2H, propyl N—C—CH₂), 2.849–3.074 (m, 4H), 3.774 (s, 3, O—CH₃), 3.623–3.880 (m, 1.5H),
 35 4.089–4.154 (m, 0.5H), 4.303–4.398 (m, 1H), 7.121–7.382 (m, 4H), 7.794 & 7.858 (two s, 1H, NH).

[α]_D²⁵=−29° (25° C., CHCl₃, c=1.0139).

Preparation of (R)-5-N-(N-Propyltrifluoromethylacetamido)-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2
 40 (1H)-one (7b):

A solution of (R)-N-Propyl-N-[3-(1-(N-methoxyaminocarbonyl)-1,2,3,4-tetrahydroquinolinyl)] trifluoromethylacetamide (6b, 5.92 g, 16.5 mmol) in chloroform (80 ml) was degassed with argon and heated to hear
 45 reflux on the steam bath. Bis(trifluoroacetoxy)iodobenzene (8.50 g, 19.8 mmol) was added portionwise. The reaction was exothermic. After the addition was complete, the solution was stirred at reflux for 5 minutes. The reaction mixture was cooled, diluted with pentane, and washed twice with
 50 saturated sodium bicarbonate solution and once with brine. The solution was dried (MgSO₄), and the solvent was removed under vacuum to leave an oil (9.2 g). Purification by flash chromatography (230–400 mesh silica gel; 30% ethyl acetate in hexane) gave the title compound (7b) as an
 55 amber oil (3.75 g, 61% yield).

NMR (CDCl₃, TMS) δ: 0.918–0.967 (m, 3H, propyl CH₃), 1.60–1.87 (m, 2H, propyl N—C—CH₂), 2.91–3.08 (m, 1H), 3.16–3.42 (m, 2H), 3.63–3.79 (m, 1H), 4.089 & 4.101 (two
 60 s, 3H, O—CH₃), 3.97–4.26 (m, 2.6H), 4.42–4.56 (m, 0.4H), 6.884–7.097 (m, 4H).

Preparation of (R)-5-Propylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (8b):

A solution of (R)-5-N-(N-Propyltrifluoromethylacetamido)-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (7b, 3.50 g, 9.79 mmol) in methanol (30 ml) and water (5 ml) was cooled in ice and potassium hydroxide (45.6% in water, 1.0 ml, 11.9

mmol) was added. The mixture was stirred at 0° C. for 1 hour and at room temperature for 18 hours. The solvent was concentrated under vacuum, and the residue was partitioned between water and diethylether. The aqueous layer was saturated with sodium chloride and extracted again with diethylether. The combined organic extract was dried (MgSO₄), and the solvent was removed under vacuum to leave an oil (2.43 g). Purification by flash chromatography (230–400 mesh silica gel; ethyl acetate to 20% tetrahydrofuran in ethyl acetate) gave the title compound (8b) as a solid (1.79 g, 69% yield).

NMR (CDCl₃, TMS) δ0.927 (t, j=7.4 Hz, 3H, propyl CH₃), 1.112 (br. s, 1H, NH), 1.507 (sextet, j=7.3 Hz, 2H, propyl N—C—CH₂), 2.629–2.804 (m, 3H), 3.062 (dd, j=4.2 & 16.0 Hz, 1H), 3.291 (heptet, j=4.0 Hz, 1H, N—CH), 3.557 (dd, j=7.7 & 12.2 Hz, 1H), 4.066–4.156 (m, 1H), 4.088 (s, 3H, O—CH₃), 6.887 (d, j=7.4 Hz, 1H), 6.961 (d, j=7.3 Hz, 1H), 7.036 (t, j=7.5 Hz, 1H).

A sample was dissolved in diethylether and excess ethereal hydrochloric acid was added. The precipitate was filtered, washed well with diethylether, and crystallized from methanol/diethylether to give the hydrochloride salt of the title compound (8b) as a yellow solid (m.p. 229° C. dec.). [α]_D²⁵ = -31° (25° C., CH₃OH, c=1.1099).

EXAMPLE 4

(R)-5-Dipropylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (9b)

Following the procedure of Example 3, a mixture of (R)-5-Propylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (8b, 0.93 g, 3.56 mmol), 1-iodopropane (3.05 g, 17.9 mmol), and potassium carbonate (1.97 g, 14.3 mmol) in acetonitrile (25 ml) was heated at reflux under nitrogen for 17 hours. 1-Iodopropane (3.05 g, 17.9 mmol) was again added, and the mixture was refluxed for an additional 5 hours. The solvent was removed under vacuum, and the mixture was diluted with water and extracted twice with diethylether. The organic extract was washed with brine and dried (MgSO₄). The solvent was removed under vacuum to leave an amber oil. Purification by flash chromatography (230–400 mesh silica gel; 10–25% ethyl acetate in hexane) gave the title compound (9b) as an amber oil (0.87 g, 90% yield).

NMR (CDCl₃, TMS) δ0.894 (t, j=7.3 Hz, 6H, propyl CH₃), 1.456 (sextet, j=7.3 Hz, 4H, propyl N—C—CH₂), 2.524 (m, 8 lines, 4H, propyl N—CH₂), 2.823–2.975 (m, 2H), 3.277 (heptet, j=5.1 Hz, 1H), 3.435 (t, j=11.4 Hz, 1H), 4.076 (s, 3H, O—CH₃), 4.168 (dd, j=4.1 & 11.5 Hz, 1H), 6.883 (d, j=7.6 Hz, 1H), 6.942 (d, j=7.5 Hz, 1H), 7.021 (t, j=7.6 Hz, 1H).

The compound was dissolved in diethylether, and excess ethereal hydrochloric acid was added. The precipitate was filtered, washed with diethylether, and crystallized from methanol/diethylether to give the hydrochloride salt of the title compound (9b) as an off-white solid (1.12 g, m.p. 175°–176° C.).

[α]_D²⁵ = -13° (25° C., CH₃OH, c=1.0682).

EXAMPLE 5

(R)-5-Methylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one (8a) (Scheme 2: Alternative Method of Example 1)

Preparation of (R)-2-(Methoxycarbonylamino)-3-phenylpropanoic acid (2):

A solution of D-phenylalanine (25.00 g, 0.151 mol) and sodium hydroxide (6.05 g, 0.151 mol) in water (170 ml) and

tetrahydrofuran (225 ml) was cooled to -15°C ., and a solution of methyl chloroformate (18.6 g, 0.197 mol) in tetrahydrofuran (50 ml) was added dropwise. When the addition was half complete, a solution of sodium hydroxide (9.10 g, 0.227 mol) in water (20 ml) was also added dropwise. When the additions were complete, the mixture was stirred at room temperature for an additional 2 hours and acid fixed with 10% hydrochloric acid. The acid was extracted twice with diethylether, and the extracts were washed with brine and dried (MgSO_4). The solvent was removed under vacuum to leave a clear oil (38.38 g). HPLC 6.01 min (78.9%), 7.32 min (5.0%), 8.29 min (10.8%), 8.42 min (0.5%), 9.95 min (1.2%), 10.27 min (2.8%), 11.92 min (0.7%).

15 Preparation of (R)-N-Methoxy-2-(methoxycarbonylamino)-3-phenylpropanamide (3):

A solution of sodium carbonate (10.20 g, 96.2 mmol) in water (170 ml) was added to a solution of the acid (-0.148 mol) in methylene chloride. Methoxylamine hydrochloride (14.2 g, 0.170 mol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (31.21 g, 0.163 mol) were added, and the mixture was stirred at room temperature for 22 hours. The mixture was diluted with tetrahydrofuran (to dissolve the precipitate) and the layers were separated. The aqueous layer was extracted with 1:1 tetrahydrofuran/diethylether, and the combined organic extracts were washed with 10% hydrochloric acid solution and saturated sodium bicarbonate solution, and the solution was dried (MgSO_4). The solvent was removed under vacuum to leave a white solid (34.2 g). Crystallization from ethyl acetate gave colorless crystals (22.6 g, m.p. $154^{\circ}\text{--}155^{\circ}\text{C}$.).

$[\alpha]_D^{+5}$ (25°C ., CH_3OH , 1.0450).

Preparation of Methyl (R)-N-(1,2,3,4-Tetrahydro-1-methoxy-2-oxo-3-quinolinyl)carbamate (4):

5 A suspension of (R)-N-Methoxy-2-(methoxycarbonylamino)-3-phenylpropanamide (11.25 g, 44.6 mmol) in 1,2-dichloroethane (170 ml) was cooled in ice and trifluoroacetic acid (9.25 ml, 13.7 g, 0.120 mol) was added. Bis(trifluoroacetoxy)iodobenzene (19.78 g, 0.046 mol) was added portionwise over 10 min at 0°C ., and the mixture was stirred at the same temperature for 1 hour. The mixture was washed with 10% sodium carbonate solution and dried (MgSO_4). The solvent was removed under vacuum to leave an amber oil (19.58 g). Purification by flash chromatography (230–400 mesh silica gel, 40–50% ethyl acetate in hexane) gave an amber oil which solidified (9.45 g, 85% yield). A sample (1.5 g) was crystallized from ethyl acetate/hexane to give white crystals (1.36 g, m.p. $117^{\circ}\text{--}119^{\circ}\text{C}$.). $[\alpha]_D^{+34}$ (25°C ., CH_3OH , 0.9279).

10 Preparation of (R)-3-Methylamino-1,2,3,4-tetrahydroquinoline maleate (5):

A solution of (R)-N-(1,2,3,4-Tetrahydro-1-methoxy-2-oxo-3-quinolinyl)carbamate (7.27 g, 29.1 mmol) in dry tetrahydrofuran (100 ml) was cooled in a water bath (-10°C .), and boron dimethylsulfide complex (10.0M, 17.5 ml, 6.0 eq) was added. The mixture was stirred in the water bath for 1 hour and at room temperature for 1.5 hours. The mixture was heated at reflux on the steam bath for 30 hours, cooled in ice, and 10% hydrochloric acid (40 ml) was added dropwise. The mixture was refluxed on the steam bath for 1.5 hours, cooled in ice, and basified with 45% potassium hydroxide. The mixture was extracted twice with diethylether, and the combined extracts were washed with brine and dried (MgSO_4). The solvent was removed under vacuum to leave a clear oil 4.89 g. The compound (0.029 mol) was combined with maleic acid (3.38 g, 0.029 mol), and the mixture was crystallized from methanol/

diethylether to give the title compound as off-white crystals (5.77 g, 71% yield, VPC shows 5.451 min (100%)). A sample (1.0 g) was recrystallized from methanol/diethylether to give a colorless solid (m.p. 173.5°–175° C.). Preparation of (R)-Methyl-(1,2,3,4-tetrahydro-3-quinolinyl) carbamic acid, phenylmethyl ester (6):

A solution of (R)-1,2,3,4-tetrahydro-N-methyl-3-quinolinamine (6.00 g, 31.5 mmol) and triethylamine (4.8 g, 47.4 mmol) in chloroform (200 ml) was cooled to –30° C., and a solution of benzyl chloroformate (5.68 g, 95% pure, 31.6 mmol) in chloroform (50 ml) was added dropwise. The mixture was stirred at –30° C. to 0° C. for 2 hours, and 10% sodium carbonate solution (100 ml) was added, the mixture was stirred at room temperature for 1 hour, diluted with pentane and diethylether, and the layers were separated. The aqueous layer was extracted with diethylether, and the combined organic extracts were washed with brine and dried (MgSO₄). The solvent was removed under vacuum to leave an oil (10.91 g). Purification by flash chromatography (230–400 mesh silica gel; 4:1 hexane/ethyl acetate) gave an oil (7.53 g, 81% yield). A sample (0.57 g) was crystallized from ethyl acetate/hexane to give white crystals (0.42 g, m.p. 78.5°–80° C.). [α]_D = –47° (25° C., CH₃OH, 0.8166). Preparation of (R)-Methyl-[1,2,3,4-tetrahydro-1-(methoxyamino)carbonyl]-3-quinolinyl]carbamic acid, phenylmethyl ester (7):

A solution of (R)-methyl-(1,2,3,4-tetrahydro-3-quinolinyl)carbamic acid, phenylmethyl ester (3.81 g, 12.86 mmol) and triethylamine (3.9 g, 39 mmol) in dry tetrahydrofuran (50 ml) was rapidly added with stirring to a solution of phosgene (1.93M in toluene) in tetrahydrofuran (100 ml) at 0° C. The cold bath was removed, and the mixture was stirred for 1.25 hours. Methoxylamine (2.15 g, 25.7 mmol) and triethylamine (7.9 g, 78 mmol) were added, and the mixture was stirred at room temperature for 3 days. The mixture was diluted with diethylether and washed with water and brine. The solution was dried (MgSO₄), and the solvent was removed under vacuum to leave an oil (5.13 g, >100%) which was sufficiently pure for the next step. A sample (0.91 g) was purified via flash chromatography (230–400 mesh silica gel; 50% ethyl acetate/hexane) to give an oil. [α]_D = +38° (25° C., CH₃OH, 0.9805), as spec. m+ at m/z 369. Strongest peaks at m/z 43, 65, 91, 118, 130, 158, 173, 204. Exact mass calcd. for C₂₀H₂₃N₃O₄: 369.1688. Found: 369.1682.

Preparation of Methyl-(1,2,5,6-tetrahydro-1-methoxy-2-oxo-4H-imidazo[4,5,1-ij]quinolinyl-5-yl)carbamic acid, phenylmethyl ester (8):

A solution of (R)-Methyl-[1,2,3,4-tetrahydro-1-(methoxyamino)carbonyl]-3-quinolinyl]carbamic acid, phenylmethyl ester (7.26 g, 19.7 mmol) in chloroform (150 ml) was degassed with argon, and the mixture was cooled to –5° C. in an ice-salt bath. Bis(trifluoroacetoxy)iodobenzene (10.14 g, 23.6 mmol) was added, and the mixture was stirred at –5° to 0° C. for 4 hours and at room temperature for 2 hours and was stored at –15° C. overnight. The mixture was washed with 10% sodium carbonate solution, and the aqueous washings were back extracted with diethylether. The combined organics were dried (MgSO₄), and the solvent was removed under vacuum to leave a brown oil (10.7 g). Purification by flash chromatography (230–400 mesh silica gel, 50% ethyl acetate in hexane) gave an amber oil which slowly solidified (5.67 g, 78%). A sample (0.54 g) was crystallized from ethyl acetate/hexane to give off-white crystals (0.41 g, m.p. 105°–106.5° C.). [α]_D = +44° (25° C., CH₃OH, 0.7311).

Preparation of (R)-5,6-Dihydro-1-methoxy-5-(methylamino)-4H-imidazo[4,5,1-ij]quinolin-2-one (8a)

Selective hydrogenation:


A mixture of methyl-(1,2,5,6-tetrahydro-1-methoxy-2-oxo-4H-imidazo[4,5,1-ij]quinolinyl-5-yl)carbamic acid phenylmethyl ester (1.48 g, 4.03 mmol) and 10% palladium on carbon (0.25 g) in absolute ethanol (100 ml) was stirred under one atmosphere of hydrogen. After taking up one equivalent of hydrogen, the mixture was filtered through celite, and the solvent was removed under vacuum to leave an oil.

Preparation of (R)-5,6-Dihydro-5-(methylamino)-4H-imidazo[4,5,1-ij]quinolin-2(1H)-one, monohydrochloride (Not a compound of the subject invention, disclosed in U.S. Pat. No. 5,273,975:

Exhaustive hydrogenation:


A mixture of methyl-(1,2,5,6-tetrahydro-1-methoxy-2-oxo-4H-imidazo[4,5,1-ij]quinolinyl-5-yl)carbamic acid phenylmethyl ester (1.48 g, 4.03 mmol) and 20% palladium hydroxide on carbon (0.50 g) in absolute ethanol (100 ml) was shaken in a Parr apparatus with an initial hydrogen pressure of 50 psi for 17 hours. The mixture was filtered through celite, and the solvent was removed under vacuum to leave an oil (0.86 g). The compound was dissolved in methanol, and excess ethereal hydrochloric acid was added. The mixture was diluted with diethylether, and the precipitate was filtered and crystallized from methanol/diethylether to give an off-white solid (0.61 g, 63% yield, m.p. 310°-311° C.). $[\alpha]_D^{25} = -30^\circ$ (25° C., CH₃OH, 1.0182).

SCHEME 1




-continued


SCHEME 1




4a. R = CH₃
4b. R = CH₂CH₂CH₃




5a. R = CH₃
5b. R = CH₂CH₂CH₃




6a. R = CH₃
6b. R = CH₂CH₂CH₃




7a. R = CH₃
7b. R = CH₂CH₂CH₃



8a. R = CH₃
8b. R = CH₂CH₂CH₃







SCHEME 2




17

-continued
SCHEME 2



What is claimed is:

1. A compound of the following structural formula:



and pharmaceutically acceptable salts thereof wherein.

R_1 and R_2 are independently hydrogen, C_{1-6} alkyl or R_1 and R_2 are joined to form pyrrolidine, piperidine, morpholine or imidazole;

X is OCH_3 , SO_2R_3 , SO_2CF_3 or CN ;

R_3 is C_{1-6} alkyl or a C_{5-10} aromatic ring (optionally substituted with a halogen, hydroxyl or C_{1-6} alkyl (optionally substituted with halogen or hydroxyl)); and

Y is hydrogen, Cl , Br , F , CN , $CONR_1R_2$, CF_3 , OCH_3 , $SO_2NR_1R_2$.

2. The compound of claim 1 wherein R_1 and R_2 are each propyl.

3. The compound of claim 1 wherein R_1 and R_2 are each methyl.

4. The compound of claim 1 wherein X is $-OCH_3$.

5. The compound of claim 1 wherein Y is hydrogen.

6. The compound of claim 1 which is

a) (R)-5-Methylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]-quinolin-2(1H)-one.

b) (R)-5-Dimethylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]-quinolin-2(1H)-one.

c) (R)-5-Propylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]-quinolin-2(1H)-one, or

d) (R)-5-Dipropylamino-1-methoxy-5,6-dihydro-4H-imidazo[4,5,1-ij]-quinolin-2(1H)-one.

7. A method for treating anxiety disorders in animal or human hosts comprising the administration of a pharmaceutically effective amount of a compound of Formula I as set forth in claim 1.

8. The method of claim 7 wherein said compound is orally administered in an amount of from about 10 mg to about 1200 mg per day.

* * * * *

- ✓ 9. (R)-3-Methylamino-1,2,3,4-tetrahydroquinoline maleate.
- ✓ 10. (R)-Methyl-(1,2,3,4-tetrahydro-3-quinoliny) carbamic acid, phenylmethyl ester.
- 5 ✓ 11. (R)-Methyl-[1,2,3,4-tetrahydro-1-[(methoxyamino)carbonyl]-3-quinoliny]carbamic acid, phenylmethyl ester.
- ✓ 12. Methyl-(1,2,5,6-tetrahydro-1-methoxy-2-oxo-4H-imidazo[4,5,1-ij]quinoliny-5-yl)carbamic acid, phenylmethyl ester.

10

DECLARATION (37 CFR 1.63) AND POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled HETEROCYCLIC AMINES HAVING CENTRAL NERVOUS SYSTEM ACTIVITY, Docket No. 4830.P CP, the specification of which

☐ is attached hereto.

☐ was filed on as Application Serial No. and was amended on .

☒ was filed on 17 June 1994 as PCT International Application No. PCT/US94/06648 and was amended under PCT Article 19 on , if applicable.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with 37 CFR 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119, of any foreign application(s) for patent or inventor's certificate or of any PCT International Application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application for patent or inventor's certificate or any PCT International Application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application on which priority is claimed:

Application Serial No.	Country	Filing Date (Day/Month/Year)	Priority Claimed (Yes/No)
---------------------------	---------	---------------------------------	------------------------------

I hereby claim the benefit under Title 35, United States Code, Section 120, of any United States application(s) or PCT International Application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application(s) in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a), which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

<u>Application Serial No.</u>	<u>Filing Date</u>	<u>Status (Patented, Pending, Abandoned)</u>
08/097,608	07/27/93	Pending

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

I hereby appoint Donald L. Corneglio (Registration 30,741), James D. Darnley, Jr. (Registration No. 33,673), Martha A. Gammill (Registration No. 31,820), William G. Jameson (Registration No. 27,199), Bruce Stein (Registration No. 27,231), Lawrence T. Welch (Registration No. 29,487), Sidney B. Williams, Jr. (Registration No. 24,949), and Thomas A. Wootton (Registration No. 35,004), all registered to practice before the Patent and Trademark Office as my attorneys with full power of substitution and revocation to prosecute this application and all divisions and continuations thereof and to transact all business in the Patent and Trademark Office connected therewith and request that all correspondence and telephone communications be directed to the following person(s) at the mailing address and telephone number hereafter given:

Name	:	Donald L. Corneglio
Registration No.	:	30,741
Address	:	Corporate Intellectual Property Law The Upjohn Company Kalamazoo, Michigan 49001
Telephone No.	:	(616) 385-4532

Inventor's Signature Arthur G. Romero Date June 30, 1994

Full name of sole or first inventor: Arthur G. Romero

Residence: Kalamazoo, MI

Citizenship: USA

Post Office Address: 2721 Lake Chevy Chase Drive
Kalamazoo, MI 49008

PL-63.1/Long Form
06/93



Pharmacia & Upjohn

January 27, 1999

Commissioner of Patents and Trademarks
Washington, DC 20231

Sir:

Pursuant to the authority delegated to me by the Board of Directors of Pharmacia & Upjohn Company, I, Jack J. Jackson, Senior Vice President, Pharmacia & Upjohn Company, hereby delegate the following:

Raymond G. Arner, Registration No. 32,958,
William G. Jameson, Registration No. 27,199,
Lori L. Kerber, Registration No. 41,113,
Ruth H. Newton, Registration No. 26,657
Bruce A. Pokras, Registration No. 32,748
Bruce Stein, Registration No. 27,231,
Thomas A. Wootton, Registration No. 35,004, and
Lucy X. Yang, Registration No. 40,259

without any approval by the Board, to sign on behalf of the Company the following papers incidental to intellectual property prosecution before the United States Patent and Trademark Office, and where required by the nature of the document executed pursuant thereto, the Secretary and any Assistant Secretary of this Company are hereby authorized to affix and attest the seal of Pharmacia & Upjohn Company on such documents:

1. Appointment and revocation of powers of attorney by Company assignee;
2. Oath or Declaration for filing by Company assignee or on behalf of the inventor when signature of the inventor is not available;
3. Assent of Company assignee to application for a reissue;
4. Assent of Company assignee to express abandonment;
5. Disclaimer or dedication by Company assignee of claims of a patent or of any terminal part of the term of a patent or application;

6. In the case of interference proceedings, to make preliminary statements on behalf of the applicant or patentee, and to consent to concessions of priority, disclaimers of invention and abandonments of contest on behalf of Company;
7. Consent of Company assignee to the addition, removal or change of inventors in patents or patent applications;
8. Declaration of Ownership;
9. Petition for Extension of Patent Term.

The signature of the delegates identified above shall constitute complete and formal execution of the document by the Company.

PHARMACIA & UPJOHN COMPANY

By: Jack J. Jackson
Jack J. Jackson

Title: Senior Vice President

Date: February 10, 1999

I hereby certify that Jack J. Jackson, who has signed the above delegation of authority, is a lawful representative of Pharmacia & Upjohn Company and is authorized to execute this document on behalf of Pharmacia & Upjohn Company, said Company having a place of business in Kalamazoo, Michigan (Address: 7000 Portage Road, Kalamazoo, Michigan 49001).

By: Robert J. Meisenhelder II
Robert J. Meisenhelder II

Title: Assistant Secretary

Date: March 1, 1999